

Table 5.A.1. -- Results

Outcome Measure	Restrictive strategy n (%)	Liberal strategy n (%)	P value
30-day mortality	78 (18.7%)	98 (23.3%)	0.11
60-day mortality	95 (22.7%)	111 (26.5%)	0.23
ICU mortality	56 (13.4%)	68 (16.2%)	0.29
Hospital mortality	93 (22.2%)	118 (28.1%)	0.05
Multiple organ dysfunction adjusted score	10.7±7.5	11.8±7.7	0.03

Table 5.A.2. -- Results

Variable	Restrictive-strategy N=320	Liberal-strategy N=317	P value
Patients transfused	146 (46%)	310 (98%)	<0.001
Total number of transfusions	301	542	<0.001
Average unit exposure per patient	0.9 ± 2.6 units	1.7 ± 2.2 units	<0.001
PICU mortality	11 (3%)	8 (3%)	0.50
28-day mortality	14 (4%)	14 (4%)	0.98
New MODS	38 (12%)	39 (12%)	NI

Table 5.A. 3. -- Results

Variable	Restrictive-strategy N=320	Liberal-strategy N=317	P value
Number of dysfunctional organs	1.6 ± 1.4 units	1.5 ± 1.2 units	0.87
Nosocomial infections	65 (20%)	79 (25%)	0.16
At least 1 adverse event	97 (30%)	90 (28%)	0.59
Mechanical ventilation	6.2 ± 5.9 days	6.0 ± 5.4 days	0.76
ICU stay after randomization	9.5 ± 7.9 days	9.9 ± 7.4 days	0.39

References (5.A.3.)

7. Istaphanous, G., Wheeler, D., Lisco, S., & Shander, A. (2011). Red blood cell transfusion in critically ill children: A narrative review. *Pediatr Crit Care Med*, 12(2):174-83. doi:10.1097/PCC.0b013e3181e30d09
10. Lacroix, J., Herbert, P.C., Hutchinson, J.S., Hume, H.A. et al. (2007). Transfusion strategies for patients in pediatric intensive care units. *N Engl J Med*, 356(16):1609-1619.
11. Hebert, P.C., Wells, G., Blajchman, M.A., Marshall, J., Martin, C., Pagliarello, G., Tweeddale, M., Schweitzer, I., Yetisir, E., & the Transfusion Requirements in Critical Care Investigators for the Canadian Critical Care Trials Group. (1999). A multicenter, randomized, controlled clinical trial of transfusion requirements in critical care. *N Engl J Med*, 340:409-17.