

Appropriate Emergency Department Blood Testing for Children with Sickle Cell Disease

Section 1. Basic Measure Information

1.A. Measure Name

Appropriate Emergency Department Blood Testing for Children with Sickle Cell Disease

1.B. Measure Number

0215

1.C. Measure Description

Please provide a non-technical description of the measure that conveys what it measures to a broad audience.

This measure assesses the percentage of children younger than 18 years of age identified as having sickle cell disease (SCD) presenting to an emergency department (ED) with fever during the measurement year, who had a pulse oximetry reading, complete blood count, reticulocyte count, and blood culture within 60 minutes following initial contact. A higher proportion indicates better performance as reflected by appropriate testing.

Approximately 2,000 infants are born with SCD in the United States each year, a condition that occurs predominantly in people of African and Hispanic descent. SCD is a chronic hematologic disorder, characterized by the presence of hemoglobin S. From infancy onward, the presence of this hemoglobin variant can lead to an array of serious medical conditions. Because children with SCD are susceptible to spleen damage, a condition that compromises their ability to deal with infection, they are at high risk for developing septicemia and meningitis from *Streptococcus pneumoniae* and other encapsulated bacteria. These illnesses can rapidly become life-threatening. Any temperature greater 38°C (100.4°F) should be immediately evaluated; this often requires a visit to the ED for timely treatment. Assessment for fever in children with SCD should include a pulse oximetry reading, complete blood count, reticulocyte count, and a blood culture. Ideally, the values collected in the ED can be compared with baseline values that should be available as part an established database profile. There are no existing quality measures for appropriate blood testing in children with SCD.

This measure uses medical record data to calculate four rates; each specified action must occur within 60 minutes following initial contact in the ED:

1. The percentage of children who had a pulse oximetry performed.
2. The percentage of children who had a complete blood count performed.
3. The percentage of children who had a reticulocyte count performed.

4. The percentage of children who had a blood culture performed.

An overall rate is calculated as the percentage of children who had a pulse oximetry reading, complete blood count, reticulocyte count, and blood culture within the same 60 minute period following initial contact in the ED.

1.D. Measure Owner

The Quality Measurement, Evaluation, Testing, Review, and Implementation Consortium (Q-METRIC).

1.E. National Quality Forum (NQF) ID (if applicable)

Not applicable.

1.F. Measure Hierarchy

Please note here if the measure is part of a measure hierarchy or is part of a measure group or composite measure. The following definitions are used by AHRQ:

- 1. Please identify the name of the collection of measures to which the measure belongs (if applicable). A collection is the highest possible level of the measure hierarchy. A collection may contain one or more sets, subsets, composites, and/or individual measures.**

This measure is part of the Q-METRIC Sickle Cell Disease Measures collection.

- 2. Please identify the name of the measure set to which the measure belongs (if applicable). A set is the second level of the hierarchy. A set may include one or more subsets, composites, and/or individual measures.**

This measure is part of the Q-METRIC Sickle Cell Disease Medical Record Data set.

- 3. Please identify the name of the subset to which the measure belongs (if applicable). A subset is the third level of the hierarchy. A subset may include one or more composites, and/or individual measures.**

Not applicable.

- 4. Please identify the name of the composite measure to which the measure belongs (if applicable). A composite is a measure with a score that is an aggregate of scores from other measures. A composite may include one or more other composites and/or individual measures. Composites may comprise component measures that can or cannot be used on their own.**

Not applicable.

1.G. Numerator Statement

The eligible population for the numerator is the number of children younger than 18 years of age identified as having SCD presenting to an ED with fever (defined as 38°C [100.4°F] or above) during the measurement year (January 1-December 31), who had (1) a pulse oximetry reading, (2) a complete blood count, (3) a reticulocyte count, and (4) a blood culture within 60 minutes following initial contact. Eligible children are restricted to those with SCD variants identified in Table 1 (see Supporting Documents), based on appropriate ICD-9 codes as documented in the medical record. The outpatient blood tests for the management of SCD are identified in Table 2 (see Supporting Documents). Four individual numerators and one overall composite of the four numerators are calculated:

1. Pulse oximetry – The number of eligible children who had a pulse oximetry reading performed within 60 minutes following initial contact in the ED.
2. Complete blood count – The number of eligible children who had a complete blood count performed within 60 minutes following initial contact in the ED.
3. Reticulocyte count – The number of eligible children who had a reticulocyte count performed within 60 minutes following initial contact in the ED.
4. Blood culture – The number of children who had a blood culture performed within 60 minutes following initial contact in the ED.
5. Overall – The number of eligible children who had pulse oximetry, a complete blood count, a reticulocyte count, and a blood culture performed within the same 60-minute period.

Evidence of a pulse oximetry reading, blood count, reticulocyte count, and blood culture is determined through medical record review of ED visits. Documentation in the medical record must include, at minimum, a note containing the time(s) at which pulse oximetry, a complete blood count, a reticulocyte count, and a blood culture were performed.

1.H. Numerator Exclusions

- Inpatient stays, outpatient visits, urgent care visits, and acute care (evaluation and management) visits with a primary care physician are excluded from the calculation.
- Children with a diagnosis in the sampled medical record indicating one of the SCD variants listed in Table 3 (see Supporting Documents) should not be included in the eligible population unless there is also a diagnosis for a sickle cell variant listed in Table 1 (see Supporting Documents).

1.I. Denominator Statement

The eligible population for the denominator is the number of children younger than 18 years of age identified as having SCD who presented to an ED with fever (defined as 38°C [100.4°F] or above) during the measurement year, (January 1-December 31) as documented in the medical record. Eligible children are restricted to those with SCD variants identified in Table 1 (see Supporting Documents), based on appropriate ICD-9 codes as documented in the medical record.

1.J. Denominator Exclusions

- Inpatient stays, outpatient visits, urgent care visits, and acute care (evaluation and management) visits with a primary care physician are excluded from the calculation.
- Children with a diagnosis in the sampled medical record indicating one of the SCD variants listed in Table 3 (see Supporting Documents) should not be included in the eligible population unless there is also a diagnosis for a sickle cell variant listed in Table 1 (see Supporting Documents).

1.K. Data Sources

Check all the data sources for which the measure is specified and tested.

Electronic health record (EHR).

If other, please list all other data sources in the field below.

Not applicable.

Section 2: Detailed Measure Specifications

Provide sufficient detail to describe how a measure would be calculated from the recommended data sources, uploading a separate document (+ Upload attachment) or a link to a URL. Examples of detailed measure specifications can be found in the CHIPRA Initial Core Set Technical Specifications Manual 2011 published by the Centers for Medicare & Medicaid Services. Although submission of formal programming code or algorithms that demonstrate how a measure would be calculated from a query of an appropriate electronic data source are not requested at this time, the availability of these resources may be a factor in determining whether a measure can be recommended for use.

Please see Supporting Documents for the technical specifications for this measure.

Section 3. Importance of the Measure

In the following sections, provide brief descriptions of how the measure meets one or more of the following criteria for measure importance (general importance, importance to Medicaid and/or CHIP, complements or enhances an existing measure). Include references related to specific points made in your narrative (not a free-form listing of citations).

3.A. Evidence for General Importance of the Measure

Provide evidence for all applicable aspects of general importance:

- **Addresses a known or suspected quality gap and/or disparity in quality (e.g., addresses a socioeconomic disparity, a racial/ethnic disparity, a disparity for Children with Special Health Care Needs (CSHCN), a disparity for limited English proficient (LEP) populations).**
- **Potential for quality improvement (i.e., there are effective approaches to reducing the quality gap or disparity in quality).**
- **Prevalence of condition among children under age 21 and/or among pregnant women.**
- **Severity of condition and burden of condition on children, family, and society (unrelated to cost).**
- **Fiscal burden of measure focus (e.g., clinical condition) on patients, families, public and private payers, or society more generally, currently and over the life span of the child.**
- **Association of measure topic with children’s future health – for example, a measure addressing childhood obesity may have implications for the subsequent development of cardiovascular diseases.**
- **The extent to which the measure is applicable to changes across developmental stages (e.g., infancy, early childhood, middle childhood, adolescence, young adulthood).**

Sickle Cell Disease Prevalence and Incidence

SCD is one of the most common genetic disorders in the United States (Kavanagh, Sprinz, Vinci, et al., 2011). The National Heart, Lung and Blood Institute (NHLBI) estimates that 2,000 infants are born with SCD in the United States each year (NHLBI, 2002). SCD affects 70,000-100,000 children and adults in the United States, predominantly those of African and Hispanic descent (Hassell, 2010).

Sickle Cell Disease Pathology and Severity

Vaso-occlusion (the sudden blockage of a blood vessel caused by the sickle shape of abnormal blood cells) is responsible for most complications of SCD, including pain episodes, sepsis, stroke, acute chest syndrome, priapism, leg ulcers, osteonecrosis, and renal insufficiency (Steinberg, 1999). In addition, SCD can have hemolytic and infectious complications that result in morbidity and mortality in children with SCD (Kavanagh, et al., 2011).

Sickle Cell Disease Burden in Daily Life

The effect of SCD on children and families is significant; severe pain episodes and hospitalizations restrict daily activities and reflect negatively on school attendance and performance, as well as on sleep and social activities (Alvim, Viana, Pires, et al., 2005; Lemanek, Ranalli, Lukens, 2009). Although medical management of SCD continues to improve over time, 196 U.S. children died from SCD-related causes between 1999 and 2002 (Yanni, Grosse, Yang, et al., 2009).

Sickle Cell Disease Cost

In a study of health care utilization among low-income children with SCD between 2004 and 2007, 27 percent of these children required inpatient hospitalization, and 39 percent used emergency care during a year. Of these children, 63 percent averaged one well-child visit per year, and 10 percent had at least one outpatient visit with a specialist (Raphael, Dietrich, Whitmire, et al., 2009). Patients with SCD use many parts of the health care system, incurring significant costs. In 2009, mean hospital charges for children with SCD and a hospital stay were \$23,000 for children with private insurance and \$18,200 for children enrolled in Medicaid (AHRQ, 2012). Kauf and colleagues estimate the lifetime cost of health care per patient with SCD to be approximately \$460,000 (Kauf, Coates, Huazhi, et al., 2009).

Outcomes of Appropriate Emergency Department Blood Testing for Children with Sickle Cell Disease

Because the spleen is often compromised at an early age in children with SCD, infection is a frequent and serious complication; the respiratory tract, in particular, serves as a common port of entry for infectious agents. Septicemia and meningitis caused by *Streptococcus pneumoniae* are a major cause of mortality in children with SCD under the age of 2 years, as this patient population experiences a 400-fold incidence of *S. pneumoniae* compared with children without SCD (NHLBI, 2002; Taylor, Moore, 2001). Other serious illnesses in children with SCD, such as acute chest syndrome, also involve fever and the presence of bacteria. Therefore, the presence of fever in children with SCD should be approached with high suspicion for systemic infection, and any febrile illness should be evaluated immediately. It is crucial that families and clinicians understand that in children with SCD, a temperature over 38.5°C is an emergency (American Academy of Pediatrics [AAP], 2002; NHLBI, 2002; Pack-Mabien, Haynes Jr, 2009). (Note that for testing purposes, the Q-METRIC quality measure drops the fever threshold to 38.0°C from the 38.5°C standard used in many clinical guidelines.)

Among the essential tests for children with SCD being evaluated for fever in the ED are a pulse oximetry reading, complete blood count, reticulocyte count, and a blood culture (NHLBI, 2002). Pulse oximetry is used to monitor arterial saturation and indicates the presence of hypoxemia (Fitzgerald, Johnson, 2001). Observed changes in values obtained from a complete blood count (which includes Q-METRIC SCD Measure 14, ED Blood Testing) may support the suspicion of infection. A reticulocyte count documents the efficiency of the marrow response and is used in determining risk of transient red cell aplasia, acute hepatic sequestration, and acute splenic sequestration complication. A blood culture probes for the presence of bacteria in the blood (NHLBI, 2002).

All children with SCD should have a set of baseline values for these test results on file so these values can be used for comparison during times of acute illness. It is important that health care providers in the ED are able to access this information quickly. Also, children with SCD should be followed at a practice or center with 24-hour access to medical consultants, hematology and microbiology laboratories, and a blood bank, among other services (NHLBI, 2002). Beyond having access to information and extended services, ED staff treating children with SCD must be skilled in providing complex care and interventions, including assessment, infection control, pain management, and appropriate understanding of complex hematological and immunological

issues. This care is often delivered in the psychosocially complicated context of chronic illness for the patient and family (Taylor, Moore, 2001).

A snapshot view of ED use by pediatric SCD patients at a major urban children's hospital found that in 97 percent of visits for fever, a complete blood count with differential and reticulocyte count were performed; 90 percent of fever visits had a blood culture sent; and hospital admissions occurred in 92 percent of patient visits for fever (Kunkel, Rackoff, Katolik, et al., 1994). Further, 53 percent of ED visits occurred at night, and 36 percent were on the weekend. In this same study, a query of families involved in the visits (40 percent contact rate) found that the mean duration of symptoms before the ED visit for fever was 0.4 days. Failure to respond to home treatment was the most common reason provided for visiting the ED (46.3 percent); worsening symptoms was cited as a reason 22.2 percent of the time. All families reported that they brought their children to the ED after initial home treatment and consultation with a hematology nurse or physician. In a study of the Healthcare Cost and Utilization Project (HCUP) inpatient and ED databases, Brousseau and colleagues found that for acute care encounters per year in children with SCD ages 1 to 9 years old, 57.4 percent had one to two visits per year, and 8.6 percent had three to 10 encounters. For children with SCD ages 10 to 17, those numbers were 51.0 percent and 12.6 percent, respectively (Brousseau, Owens, Mosso, et al., 2010).

This measure assesses whether children younger than 18 years of age with SCD presenting to an ED with a fever received appropriate blood testing within 60 minutes of initial contact: a pulse oximetry reading and orders for a complete blood count, reticulocyte count, and a blood culture. The measure does not change across developmental stages.

Performance Gap

A common complaint among patients with SCD and their families is that of receiving inappropriate care in the ED. The issue is not so much a lack of knowledge about the disease on the part of ED staff, but rather that staff are unfamiliar with the individual presenting. Given the fast pace of most EDs and the needs of the young patient with SCD for rapid and efficient evaluation, quick access to individual care plans and patient records (including values for baseline blood work) as described above is essential. Electronic health records (EHRs), patient information cards, and phone calls to patient providers are all means to transmit information quickly during an emergency (NHLBI, 2002).

Underdeveloped quality standards for treating children with SCD presenting to the ED is another identified gap (Tanabe, Dias, Gorman, 2013). Caring for children in the ED is complex from both a medical and behavioral perspective. Children vary developmentally; their ability to communicate and degree of independence may vary widely. For children with SCD, physical challenges and neurocognitive deficits can make ED visits even more complicated. The quality measures developed by Q-METRIC are one approach to rectifying this problem, as are quality of care indicators recently proposed (Wang, Kavanagh, Litle, et al., 2011).

3.B. Evidence for Importance of the Measure to Medicaid and/or CHIP

Comment on any specific features of this measure important to Medicaid and/or CHIP that are in addition to the evidence of importance described above, including the following:

- **The extent to which the measure is understood to be sensitive to changes in Medicaid or CHIP (e.g., policy changes, quality improvement strategies).**
- **Relevance to the Early and Periodic Screening, Diagnostic and Treatment benefit in Medicaid (EPSDT).**
- **Any other specific relevance to Medicaid/CHIP (please specify).**

Sickle Cell Disease and Medicaid/CHIP

The majority of children with SCD are enrolled in Medicaid. In 2009, 67 percent of pediatric SCD patients discharged from the hospital were enrolled in Medicaid; only 25 percent had private insurance (AHRQ, 2012). In a study of the HCUP inpatient and ED databases, Brousseau and colleagues (2010) found that for acute care encounters (ED and hospitalization) for patients with SCD, children ages 1-9 years with public insurance had 1.6 visits per year compared with 1.39 for those with private insurance and 1.10 for the uninsured.

A study of patients (most of whom were black), including children, with SCD enrolled in TennCare, Tennessee's Medicaid managed health care program, from January 1995 to December 2002 showed much higher rates of ED use compared with black patients in TennCare without SCD. For children younger than 5 years of age, the rate ratio (RR) of ED visits per 1,000 person years was 1.8 for boys (95 percent confidence interval [CI] = 1.7 to 1.9) and 2.0 for girls (1.9 - 2.2). For those ages 5 to 9 years, the RR for boys was 2.7 (2.5-2.9), and for girls it was 3.0 (2.8-3.2). For those ages 10 to 19 years, the RR for boys was 3.7 (3.5-3.9), and for girls it was 3.7 (3.4-3.7) (Shankar, Arbogast, Mitchel, et al., 2005).

Medicaid enrollment often serves as a marker of poverty. The large number of children with SCD on Medicaid suggests some of these patients may be receiving suboptimum treatment because of unstable living situations. They may not be receiving prophylactic antibiotics to help prevent bacterial infections, and they may experience delays in being taken for medical care if family situations are such that work responsibilities, school commitments for siblings, or lack of transportation make seeking prompt medical attention difficult (Tanabe, et al., 2013). Having consistent standards of care to treat these children quickly and effectively when they present in the ED is an important measure to help rectify disadvantages they face because of socioeconomic status.

3.C. Relationship to Other Measures (if any)

Describe, if known, how this measure complements or improves on an existing measure in this topic area for the child or adult population, or if it is intended to fill a specific gap in an existing measure category or topic. For example, the proposed measure may enhance an existing measure in the initial core set, it may lower the age range for an existing adult-focused measure, or it may fill a gap in measurement (e.g., for asthma care quality, inpatient care measures).

Currently, there are no quality measures for the diagnosis, assessment, or treatment of pediatric SCD.

Section 4. Measure Categories

CHIPRA legislation requires that measures in the initial and improved core set, taken together, cover all settings, services, and topics of health care relevant to children. Moreover, the legislation requires the core set to address the needs of children across all ages, including services to promote healthy birth. Regardless of the eventual use of the measure, we are interested in knowing all settings, services, measure topics, and populations that this measure addresses. These categories are not exclusive of one another, so please indicate "Yes" to all that apply.

Does the measure address this category?

- a. Care Setting – ambulatory: Yes.
- b. Care Setting – inpatient: No.
- c. Care Setting – other – please specify: No.
- d. Service – preventive health, including services to promote healthy birth: No.
- e. Service – care for acute conditions: Yes.
- f. Service – care for children with acute conditions: Yes.
- g. Service – other (please specify): No.
- h. Measure Topic – duration of enrollment: No.
- i. Measure Topic – clinical quality: No.
- j. Measure Topic – patient safety: Yes.
- k. Measure Topic – family experience with care: No.
- l. Measure Topic – care in the most integrated setting: No.
- m. Measure Topic other (please specify): Not applicable.
- n. Population – pregnant women: Not applicable.
- o. Population – neonates (28 days after birth) (specify age range): Yes; birth to 28 days.
- p. Population – infants (29 days to 1 year) (specify age range): Yes; 29 days to 1 year.
- q. Population – pre-school age children (1 year through 5 years) (specify age range): Yes; all ages in this range.
- r. Population – school-aged children (6 years through 10 years) (specify age range): Yes; all ages in this range.
- s. Population – adolescents (11 years through 20 years) (specify age range): Yes; adolescents 11 through 17 years.
- t. Population – other (specify age range): Not applicable.
- u. Other category (please specify): Not applicable.

Section 5. Evidence or Other Justification for the Focus of the Measure

The evidence base for the focus of the measures will be made explicit and transparent as part of the public release of CHIPRA deliberations; thus, it is critical for submitters to specify the scientific evidence or other basis for the focus of the measure in the following sections.

5.A. Research Evidence

Research evidence should include a brief description of the evidence base for valid relationship(s) among the structure, process, and/or outcome of health care that is the focus of the measure. For example, evidence exists for the relationship between immunizing a child or adolescent (process of care) and improved outcomes for the child and the public. If sufficient evidence existed for the use of immunization registries in practice or at the State level and the provision of immunizations to children and adolescents, such evidence would support the focus of a measure on immunization registries (a structural measure).

Describe the nature of the evidence, including study design, and provide relevant citations for statements made. Evidence may include rigorous systematic reviews of research literature and high-quality research studies.

This measure focuses on a clinical process (appropriate emergency department blood testing for children with SCD), that, if followed, results in a desirable clinical outcome (timely discovery and treatment of infection in children with SCD). The measure highlights where providers or hospitals are falling short in offering quality health care in the ED for children with SCD.

In the ED, fever in children with SCD is a very high-triage priority because the risk of sepsis is life threatening. Early identification and management of infection are critical (Tanabe et al., 2013). To that end, performing standard-of-care blood tests quickly in the ED is essential to establish a diagnosis and initiate treatment. Overall, clinical guidelines indicate that providers should perform pulse oximetry, a complete blood count, reticulocyte count, and blood culture soon after presenting at the ED. Table 4 summarizes several key sources of evidence for this measure, using the US Preventive Services Task Force (USPSTF) rankings (criteria denoted in Table 4).

5.B. Clinical or Other Rationale Supporting the Focus of the Measure (optional)

Provide documentation of the clinical or other rationale for the focus of this measure, including citations as appropriate and available.

Children with SCD are at high risk of developing bacterial infections because of the loss of splenic function by the age of 2 to 3 months. Prophylactic use of antibiotics in children until age 5 (Gaston et al., 1986) has decreased the incidence of sepsis and meningitis; routine flu vaccinations and the introduction of the 7-valent pneumococcal conjugate vaccine may also contributed to a reduced incidence of bacteremia, though the connection is still debated (Adamkiewicz et al., 2003). Still, given the susceptibility of these children to infection, a high degree of suspicion should be maintained for children with SCD who present with fever. A study by Bansil et al. reported that during a 10-year period, 16 percent of febrile children with SCD discharged from the pediatric ED of a major medical center had a serious bacterial infection: pneumonia was diagnosed most often at 13.8 percent, followed by bacteremia and urinary tract infections, both at 1.1 percent. The authors suggested that because children with SCD have lower immunity, they may be susceptible to *S. pneumoniae* serotypes not covered by the PCV (Bansil et al., 2013).

Fever is also associated with other serious SCD complications, such as acute chest syndrome, a common and life-threatening condition in children with SCD that can also be caused by bacterial infections. Acute chest syndrome is difficult to distinguish from pneumonia because both illnesses present with fever and cough. In children with SCD younger than 2 years of age, 97 percent of those with acute chest syndrome are febrile; 17.4 percent of febrile children with SCD have acute chest syndrome (Chang et al., 2013).

SCD is a chronic hematologic disorder, characterized by the presence of hemoglobin S. Oxygenated HbS functions normally but in decreased oxygen states (hypoxemia), HbS becomes distorted to the sickle shape, which leads to vaso-occlusion and ischemia. A cycle of hypoxemia and sickling is set up, which leads to SCD complications (Fitzgerald et al., 2001). Blood testing provides important information about a patient's status regarding many SCD-related conditions, including anemia, infection and fever, stroke, acute chest syndrome, and pain. Expert consensus indicates that the four tests blood tests listed below should be conducted immediately when febrile children with SCD present in the ED:

Pulse oximetry is a simple, accurate, noninvasive technique to monitor the amount of oxygen in the blood (oxygen saturation). The test is administered using a small probe that's placed on a thin part of a patient's body, such as a finger or ear lobe. The probe transmits two wavelengths of light through the vascular tissue and measures the differential absorption of the light by oxyhemoglobin and deoxyhemoglobin, allowing health care providers to assess the amount of oxygen in the blood (Johns Hopkins Medicine).

A complete blood count (CBC) provides information about the kinds and numbers of cells in the blood, especially red blood cells, white blood cells, and platelets. The test helps to check symptoms and diagnose conditions. A CBC usually includes white blood cell (WBC, leukocyte) count; white blood cell types (WBC differential); red blood cell (RBC) count; hematocrit (HCT, packed cell volume, PVC); hemoglobin (Hgb); red blood cell indices; platelet (thrombocyte) count; mean platelet volume (MPV) (WebMD, 2012).

A reticulocyte count measures the number of immature red blood cells (a.k.a., reticulocytes) released into the blood by the bone marrow. Usually reticulocytes are in the blood for about two days before developing into mature red blood cells; normally about 1 percent to 2 percent of red blood cells are reticulocytes. Reticulocyte counts rise when red blood cells are destroyed prematurely (WebMD, 2012).

A blood culture tests for infection in the blood by detecting the presence of bacteria or fungi. A bacterial infection in the blood can be serious because the blood can spread bacteria to any part of the body. To test for infection, blood is collected and placed in a cup with substances that allow bacteria or fungus to grow. If this occurs, the bacteria or fungus are checked visually by a microscope and chemically. Further sensitivity testing is conducted to determine the best antibiotic to use to kill the bacteria. Two or three samples may be taken from different veins to make sure infection is not missed. If nothing grows, the blood culture is called negative (WebMD, 2012).

A prospective study at the Children's Hospital of Philadelphia looked at the use of blood cultures in determining the presence of bacteremia in children with SCD. Because management varies for evaluating febrile illness in this population, establishing an average time to a positive culture in a continuously monitored blood culture instrument will allow for guidelines to be developed to stratify the length of time a child should be observed, based on clinical, laboratory, and social features (Norris et al., 2003).

Section 6. Scientific Soundness of the Measure

Explain the methods used to determine the scientific soundness of the measure itself. Include results of all tests of validity and reliability, including description(s) of the study sample(s) and methods used to arrive at the results. Note how characteristics of other data systems, data sources, or eligible populations may affect reliability and validity.

6.A. Reliability

Reliability of the measure is the extent to which the measure results are reproducible when conditions remain the same. The method for establishing the reliability of a measure will depend on the type of measure, data source, and other factors.

Explain your rationale for selecting the methods you have chosen, show how you used the methods chosen, and provide information on the results (e.g., the Kappa statistic). Provide appropriate citations to justify methods.

This measure is based on medical record data. Reliability testing is described below.

Data and Methods

Our testing data consisted of an audit of medical records from the three largest centers serving SCD patients in Michigan during 2012: Children's Hospital of Michigan (CHM, Detroit), Hurley Medical Center (Hurley, Flint), and the University of Michigan Health System (UMHS, Ann Arbor). Combined, these sites treat the majority of children with SCD in Michigan. Medical records for all children with SCD meeting the measure specification criteria during the measurement year were abstracted at each site. Abstracting was conducted in two phases; during Phase 1, 435 records were abstracted among the three sites. In Phase 2, an additional 237 cases were abstracted at one site. In total, 672 unique records were reviewed for children with SCD to test this measure.

Reliability of medical record data was determined through re-abstraction of patient record data to calculate the inter-rater reliability (IRR) between abstractors. Broadly, IRR is the extent to which the abstracted information is collected in a consistent manner. Low IRR may be a sign of poorly executed abstraction procedures, such as ambiguous wording in the data collection tool, inadequate abstractor training, or abstractor fatigue. For this measure, the medical record data collected by two nurse abstractors were compared.

Measuring IRR at the beginning of the abstraction is imperative to identify any misinterpretations early on. It is also important to assess IRR throughout the abstraction process

to ensure that the collected data maintain high reliability standards. Therefore, the IRR was evaluated during Phase 1 at each site to address any reliability issues before beginning data abstraction at the next site.

IRR was determined by calculating both percent agreement and Kappa statistics. While abstraction was still being conducted at each site, IRR assessments were conducted for 5 percent of the total set of unique patient records that were abstracted during Phase 1 of data collection. Two abstractors reviewed the same medical records; findings from these abstractions were then compared, and a list of discrepancies was created.

Three separate IRR meetings were conducted, all of which included a review of multiple SCD measures that were being evaluated. Because of eligibility criteria, not all patients were eligible for all measures. Therefore, records for IRR were not chosen completely at random; rather, records were selected to maximize the number of measures assessed for IRR at each site.

Results

For each of the measure numerators, 14 of 435 unique patient records (3 percent) from Phase 1 of the abstraction process were assessed for IRR across the three testing sites. Additionally, in order for a record to be abstracted for this measure, patients must meet a specific medical criterion (fever). Therefore, IRR was also assessed for this eligibility criterion. For fever, 29 of 435 unique patient records (7 percent) from Phase 1 of the abstraction process were assessed for IRR across the three testing sites.

Table 5 shows the percent agreement and Kappa statistic for each numerator and the fever eligibility criterion of this measure for each site and across all sites. The overall agreement for three of the numerators (complete blood count, reticulocyte, and blood culture) was 100 percent and the Kappa was 1.00, indicating a perfect IRR level was achieved. The overall agreement for the pulse oximetry numerator was 64 percent with a Kappa statistic of 0.10. The overall agreement for fever, an eligibility criterion, was 90 percent and the Kappa was 0.79.

Discrepancies

When discrepancies between abstractors were found, the abstractors and a study team member reopened the electronic medical record to review each abstractor's response and determine the correct answer. After discussion, a consensus result was obtained and inconsistent records were corrected for the final dataset. When consistent differences were noted between the abstractors, clarification was provided and the abstraction tool modified, where appropriate.

For the fever eligibility criterion, 26 of 29 records agreed, resulting in a 90 percent agreement and a Kappa score of 0.79. In two of the three records where there was disagreement, it was because the first temperature taken in the ED was below the fever threshold. Therefore, one of the abstractors recorded that there was not a fever. The second abstractor used a temperature reading taken later in the visit that was above the threshold to indicate that there was a fever. During the review meeting, it was clarified that the patient had to present to the ED with a fever, therefore the first temperature recorded should be used to determine eligibility. This text was also added to the abstraction tool.

The numerator that assesses pulse oximetry had a percent agreement of 64 percent and a Kappa statistic of 0.10 due to five of 14 records in disagreement. All five of the disagreements occurred in records at one hospital. While reviewing the medical records, it was discovered that pulse oximetry readings were not always listed with other vital signs. It was not initially known that the abstractor may have to navigate to another section of the record to find this information. Following review, the abstractors verified that they knew where to find pulse oximetry information, thus resolving this issue.

6.B. Validity

Validity of the measure is the extent to which the measure meaningfully represents the concept being evaluated. The method for establishing the validity of a measure will depend on the type of measure, data source, and other factors.

Explain your rationale for selecting the methods you have chosen, show how you used the methods chosen, and provide information on the results (e.g., R2 for concurrent validity).

The validity of this measure was determined from two perspectives: face validity and validity of medical record data.

Face Validity

Face validity is the degree to which the measure construct characterizes the concept being assessed. The face validity of this measure was established by a national panel of experts and advocates for families of children with SCD convened by Q-METRIC. The Q-METRIC expert panel included nationally recognized experts in SCD, representing hematology, pediatrics, and SCD family advocacy. In addition, measure validity was considered by experts in state Medicaid program operations, health plan quality measurement, health informatics, and health care quality measurement. In total, the Q-METRIC SCD panel included 14 experts, providing a comprehensive perspective on SCD management and the measurement of quality metrics for states and health plans.

The Q-METRIC expert panel concluded that this measure has a high degree of face validity through a detailed review of concepts and metrics considered to be essential to effective SCD management and treatment. Concepts and draft measures were rated by this group for their relative importance. This measure was highly rated, receiving an average score of 8.1 (with 9 as the highest possible score).

Validity of Abstracted Data

This measure was tested using medical record data. This source is considered the gold standard for clinical information; our findings indicate that these data have a high degree of face validity. This measure was tested among a total of 123 children younger than 18 years of age with sickle cell disease (Table 6). Overall, appropriate blood testing was conducted within 60 minutes of initial contact in the ED for 10 percent of children with SCD (range among the three hospitals: 7 percent-10 percent). Pulse oximetry was conducted within 60 minutes of initial contact in the ED on 51 percent of children (range: 39 percent- 100 percent). Similarly, a complete blood count was conducted within 60 minutes of initial contact in the ED for 25 percent of children (range:

20 percent-26 percent); a reticulocyte count was conducted within 60 minutes of initial contact in the ED for 24 percent of children (range: 20-25 percent); a blood culture was conducted within 60 minutes of initial contact in the ED for 39 percent of children (range: 7 percent-44 percent).

Section 7. Identification of Disparities

CHIPRA requires that quality measures be able to identify disparities by race, ethnicity, socioeconomic status, and special health care needs. Thus, we strongly encourage nominators to have tested measures in diverse populations. Such testing provides evidence for assessing measure’s performance for disparities identification. In the sections below, describe the results of efforts to demonstrate the capacity of this measure to produce results that can be stratified by the characteristics noted and retain the scientific soundness (reliability and validity) within and across the relevant subgroups.

7.A. Race/Ethnicity

The measure was tested using medical records from the three largest centers serving SCD patients in Michigan during 2012: Children’s Hospital of Michigan, Hurley Medical Center, and the University of Michigan Health System. Combined, these centers serve the vast majority of SCD patients in Michigan. While race and ethnicity data were not abstracted as part of the medical record review process, information is available from the state of Michigan for its entire population of births with an initial newborn screening result indicating SCD from 2004 to 2008. Table 7 summarizes the distribution across race and ethnicity groups for all SCD births in Michigan during that time period.

7.B. Special Health Care Needs

The medical records abstracted for this study do not include indicators of special health care needs.

7.C. Socioeconomic Status

The medical records abstracted for this study do not include indicators of socioeconomic status.

7.D. Rurality/Urbanicity

The medical records abstracted for this study do not include indicators of urban/rural residence.

7.E. Limited English Proficiency (LEP) Populations

The medical records abstracted for this study do not include indicators of LEP.

Section 8. Feasibility

Feasibility is the extent to which the data required for the measure are readily available, retrievable without undue burden, and can be implemented for performance measurement. Using the following sections, explain the methods used to determine the feasibility of implementing the measure.

8.A. Data Availability

1. What is the availability of data in existing data systems? How readily are the data available?

This measure is based on review of medical record data. The medical chart audit included records from the three largest centers serving SCD patients in Michigan during 2012: Children's Hospital of Michigan, Hurley Medical Center, and the University of Michigan Health System. Data were abstracted from EHRs at all three sites.

Medical records for 100 percent of children with SCD meeting the measure specification criteria during the measurement year were abstracted from each hospital. In total, 672 unique records were reviewed; 123 records (18 percent) met denominator criteria for this measure.

Based on the abstracted chart data, rates were calculated for each of the five numerators (see Table 6 in the Validity section above) as follows:

1. Pulse Oximetry: The percentage of children with SCD who had a pulse oximetry reading within 60 minutes following initial contact in the ED (51 percent). Pulse oximetry numerator (63) divided by denominator (123).
2. Complete Blood Count: The percentage of children with SCD who had a complete blood count within 60 minutes following initial contact in the ED (25 percent). Complete blood count numerator (31) divided by denominator (123).
3. Reticulocyte: The percentage of children with SCD who had a reticulocyte count within 60 minutes following initial contact in the ED (24 percent). Reticulocyte numerator (30) divided by denominator 123).
4. Blood Culture: The percentage of children with SCD who had a blood culture count within 60 minutes following initial contact in the ED (39 percent). Blood culture numerator (48) divided by denominator (123).
5. Overall: The overall rate is the percentage of children with SCD who had a pulse oximetry, complete blood count, reticulocyte count, and blood culture performed within 60 minutes following initial contact in the ED (10 percent). Overall numerator (12) divided by denominator (123).

Medical record abstraction for this measure was accomplished with a data collection tool developed using LimeSurvey software (version 1.92, formerly PHPSurveyor). LimeSurvey is an open-source online application based in MySQL that enables users to develop and publish surveys, as well as collect responses. The tool was piloted to determine its usability and revised

as necessary. The technical specification for this measure also underwent revisions following pilot testing.

Data abstraction was completed by experienced nurse abstractors who had undergone training for each medical record system used. Abstractors participated in onsite training during which the measure was discussed in length to include the description, calculation, definitions, eligible population specification, and exclusions. Following training, abstractors were provided with a coded list of potentially eligible cases from each of the sites. To abstract all pertinent data, two nurse abstractors reviewed the electronic records. In addition to the specific data values required for this measure, key patient characteristics, such as date of birth and hemoglobin variant type, were also collected.

Abstraction Times

In addition to calculating IRR, the study team assessed how burdensome it was to locate and record the information used to test this measure by having abstractors note the time it took to complete each record. During Phase 1, On average, the abstractors spent 12 minutes per eligible SCD case abstracting the data for this measure, with times ranging from 5-45 minutes.

2. If data are not available in existing data systems or would be better collected from future data systems, what is the potential for modifying current data systems or creating new data systems to enhance the feasibility of the measure and facilitate implementation?

The proposed measure was determined to be feasible by Q-METRIC using electronic health record data from the three largest centers serving SCD patients in Michigan during 2012.

8.B. Lessons from Use of the Measure

1. Describe the extent to which the measure has been used or is in use, including the types of settings in which it has been used, and purposes for which it has been used.

To our knowledge, this measure is not currently in use anywhere in the United States.

2. If the measure has been used or is in use, what methods, if any, have already been used to collect data for this measure?

Not applicable.

3. What lessons are available from the current or prior use of the measure?

Not applicable.

Section 9. Levels of Aggregation

CHIPRA states that data used in quality measures must be collected and reported in a standard format that permits comparison (at minimum) at State, health plan, and provider levels. Use the following table to provide information about this measure's use for reporting at the levels of aggregation in the table.

For the purpose of this section, please refer to the definitions for provider, practice site, medical group, and network in the Glossary of Terms.

If there is no information about whether the measure could be meaningfully reported at a specific level of aggregation, please write "Not available" in the text field before progressing to the next section.

Level of aggregation (Unit) for reporting on the quality of care for children covered by pp

State level Can compare States*

Intended use: Is measure intended to support meaningful comparisons at this level? (Yes/No)

No.

Data Sources: Are data sources available to support reporting at this level?

No.

Sample Size: What is the typical sample size available for each unit at this level? What proportion of units at this level of aggregation can achieve an acceptable minimum sample size?

Not applicable.

In Use: Have measure results been reported at this level previously?

No.

Reliability & Validity: Is there published evidence about the reliability and validity of the measure when reported at this level of aggregation?

No.

Unintended consequences: What are the potential unintended consequences of reporting at this level of aggregation?

Not applicable.

Other geographic level: Can compare other geographic regions (e.g., MSA, HRR)

Intended use: Is measure intended to support meaningful comparisons at this level? (Yes/No)

No.

Data Sources: Are data sources available to support reporting at this level?

No.

Sample Size: What is the typical sample size available for each unit at this level? What proportion of units at this level of aggregation can achieve an acceptable minimum sample size?

Not applicable.

In Use: Have measure results been reported at this level previously?

No.

Reliability & Validity: Is there published evidence about the reliability and validity of the measure when reported at this level of aggregation?

No.

Unintended consequences: What are the potential unintended consequences of reporting at this level of aggregation?

Not applicable.

Medicaid or CHIP Payment model: Can compare payment models (e.g., managed care, primary care case management, FFS, and other models)

Intended use: Is measure intended to support meaningful comparisons at this level?

(Yes/No)

No.

Data Sources: Are data sources available to support reporting at this level?

No.

Sample Size: What is the typical sample size available for each unit at this level? What proportion of units at this level of aggregation can achieve an acceptable minimum sample size?

Not applicable.

In Use: Have measure results been reported at this level previously?

No.

Reliability & Validity: Is there published evidence about the reliability and validity of the measure when reported at this level of aggregation?

No.

Unintended consequences: What are the potential unintended consequences of reporting at this level of aggregation?

Not applicable.

Health plan*: Can compare quality of care among health plans.

Intended use: Is measure intended to support meaningful comparisons at this level?

(Yes/No)

No.

Data Sources: Are data sources available to support reporting at this level?

No.

Sample Size: What is the typical sample size available for each unit at this level? What proportion of units at this level of aggregation can achieve an acceptable minimum sample size?

Not applicable.

In Use: Have measure results been reported at this level previously?

No.

Reliability & Validity: Is there published evidence about the reliability and validity of the measure when reported at this level of aggregation?

No.

Unintended consequences: What are the potential unintended consequences of reporting at this level of aggregation?

Not applicable.

Provider Level

Individual practitioner: Can compare individual health care professionals

Intended use: Is measure intended to support meaningful comparisons at this level?
(Yes/No)

No.

Data Sources: Are data sources available to support reporting at this level?

No.

Sample Size: What is the typical sample size available for each unit at this level? What proportion of units at this level of aggregation can achieve an acceptable minimum sample size?

Not applicable.

In Use: Have measure results been reported at this level previously?

No.

Reliability & Validity: Is there published evidence about the reliability and validity of the measure when reported at this level of aggregation?

No.

Unintended consequences: What are the potential unintended consequences of reporting at this level of aggregation?

Not applicable.

Provider Level

Hospital: Can compare hospitals

Intended use: Is measure intended to support meaningful comparisons at this level?
(Yes/No)

Yes.

Data Sources: Are data sources available to support reporting at this level?

Yes.

Sample Size: What is the typical sample size available for each unit at this level? What proportion of units at this level of aggregation can achieve an acceptable minimum sample size?

The sample would comprise all children with clinical documentation of sickle cell disease presenting to the ED.

In Use: Have measure results been reported at this level previously?

No.

Reliability & Validity: Is there published evidence about the reliability and validity of the measure when reported at this level of aggregation?

No.

Unintended consequences: What are the potential unintended consequences of reporting at this level of aggregation?

Not applicable.

Provider Level

Practice, group, or facility: Can compare: (i) practice sites; (ii) medical or other professional groups; or (iii) integrated or other delivery networks**

Intended use: Is measure intended to support meaningful comparisons at this level?

(Yes/No)

No.

Data Sources: Are data sources available to support reporting at this level?

No.

Sample Size: What is the typical sample size available for each unit at this level? What proportion of units at this level of aggregation can achieve an acceptable minimum sample size?

Not applicable.

In Use: Have measure results been reported at this level previously?

No.

Reliability & Validity: Is there published evidence about the reliability and validity of the measure when reported at this level of aggregation?

No.

Unintended consequences: What are the potential unintended consequences of reporting at this level of aggregation?

Not applicable.

Section 10. Understandability

CHIPRA states that the core set should allow purchasers, families, and health care providers to understand the quality of care for children. Please describe the usefulness of this measure toward achieving this goal. Describe efforts to assess the understandability of this measure (e.g., focus group testing with stakeholders).

This measure provides families with a straightforward measure to assess how well basic levels of comprehensive care are being provided for children with SCD. Low rates for the provision of care in the emergency department are easily understood to be unsatisfactory. The simplicity of the measure likewise makes it a straightforward guide for providers and purchasers to assess how well comprehensive care, including treatment in the ED, is managed in children with SCD.

This measure has not been assessed for comprehension. The primary information needed for this measure comes from medical records data and includes basic demographics, diagnostic codes, and procedure codes, all of which are widely available. The nurse abstractors testing the measure provided feedback to refine the abstraction tool and thus the specifications. These changes are reflected in the final documentation.

Section 11. Health Information Technology

Please respond to the following questions in terms of any health information technology (health IT) that has been or could be incorporated into the measure calculation.

11.A. Health IT Enhancement

Please describe how health IT may enhance the use of this measure.

Health IT may enhance the use of this measure by providing the vehicle for ensuring timely completion of these activities, and by providing queues for these activities that are aligned with roles. For example, when a patient arrives to an ED that has performed poorly on these measures, the source of poor performance may be related to waiting times. Health IT in the triage area could trigger different decision-making that allows these patients to be seen more quickly. Another source might be documentation of completed tasks, which can be either automated by health IT or augmented by tools such as mobile entry tools for nursing staff.

11.B. Health IT Testing

Has the measure been tested as part of an electronic health record (EHR) or other health IT system?

Yes.

If so, in what health IT system was it tested and what were the results of testing?

This measure was tested using EHR review conducted at three major SCD treatment facilities in Michigan.

11.C. Health IT Workflow

Please describe how the information needed to calculate the measure may be captured as part of routine clinical or administrative workflow.

This information is most typically captured in orders or results within the EHR or computerized physician order entry (CPOE) system. It will be captured by nurses, technicians, or physicians, depending on the workflow of the clinic.

11.D. Health IT Standards

Are the data elements in this measure supported explicitly by the Office of the National Coordinator for Health IT Standards and Certification criteria (see healthit.hhs.gov/portal/server.pt/community/healthit_hhs_gov__standards_ifr/1195)?

Yes.

If yes, please describe.

The ONC's Health IT Standards explicitly address the receipt of laboratory results and other diagnostic tests into EHRs, which are directly relevant to this measure. In addition, these standards indicate the requirement for EHRs to track specific patient conditions, such as SCD. The ONC standards include the following specific requirements in the Certification criteria (Federal Register 2010) pertaining to Stage 2 Meaningful Use requirements include:

Stage 2 (beginning in 2013): CMS has proposed that its goals for the Stage 2 meaningful use criteria expand upon the Stage 1 criteria to encourage the use of health IT for continuous quality improvement at the point of care. In addition, the exchange of information in the most structured format possible is encouraged. This can be accomplished through mechanisms such as the electronic transmission of orders entered using computerized provider order entry (CPOE) and the electronic transmission of diagnostic test results. Electronic transmission of diagnostic test results include a broad array of data important to quality measurement, such as blood tests, microbiology, urinalysis, pathology tests, radiology, cardiac imaging, nuclear medicine tests, and pulmonary function tests.

Incorporate clinical lab test results into the HER as structured data:

1. Electronically receive clinical laboratory results in a structured format and display such results in human readable format.
2. Electronically display in human readable format any clinical laboratory tests that have been received with LOINC[®] codes.

3. Electronically display all the information for a test report specified at 42 CFR 493.1291(c)(1) through (7).

Generate lists of patients by specific conditions to use for quality improvement reduction of disparities outreach:

4. Enable a user to electronically update a patient's record based upon received laboratory test results. Enable a user to electronically select, sort, retrieve, and output a list of patients and patients' clinical information, based on user-defined demographic data, medication list, and specific conditions.

11.E. Health IT Calculation

Please assess the likelihood that missing or ambiguous information will lead to calculation errors.

Missing or ambiguous information in the following areas could lead to missed cases or calculation errors:

1. Child's date of birth.
2. ICD-9-CM codes selected to indicate sickle cell anemia/SCD.
3. Date and time of treatment.
4. Type of tests administered.
5. Date of tests performed.
6. Care setting.

11.F. Health IT Other Functions

If the measure is implemented in an EHR or other health IT system, how might implementation of other health IT functions (e.g., computerized decision support systems in an EHR) enhance performance characteristics on the measure?

Being able to show these measure results in health IT, especially to patients, might be transformative. Imagine, for example, an electronic white board in the room that describes "Our goals for your care" and has green, yellow, and red lights next to each of these measures. This system would be hypothesized to improve delivery of this care. Another approach that has been demonstrated to significantly improve quality is through the use of a process control system that health care administrators or leaders could monitor to ensure 100 percent compliance with these measures, employing the same types of warnings to prompt action before the time window has expired.

Section 12. Limitations of the Measure

Describe any limitations of the measure related to the attributes included in this CPCF (i.e., availability of measure specifications, importance of the measure, evidence for the focus of the measure, scientific soundness of the measure, identification of disparities, feasibility, levels of aggregation, understandability, health information technology).

This measure assesses the percentage of children younger than 18 years of age identified as having sickle cell disease (SCD) presenting to an emergency department (ED) with fever during the measurement year, who 1) received a pulse oximetry reading and 2) had a complete blood count, 3) reticulocyte count and 4) blood culture within 60 minutes following initial contact.

This measure is implemented with medical record data, and was tested with electronic medical records. The primary information needed for this measure includes date of birth, diagnosis codes, and procedure codes and dates. These data are available, although obtaining them may require a restricted-use data agreement. It also required the development of an abstraction tool and the use of qualified nurse abstractors. Continuing advances in the development and implementation of electronic medical records may establish the feasibility of regularly implementing this measure with data supplied by electronic medical records.

In future implementations, there are several considerations that may further strengthen this measure and potentially ease the burden of data collection. Specific feedback from our medical record abstractors suggested that either the time the test was ordered or the time the result was recorded in lieu of the time the test was performed would be easier to find in the medical record. The abstractors also suggested that future versions of the specification note whether any temperature above the threshold (38°C / 100.4° F) can be used to indicate fever or if abstractors should use only the first temperature recorded to determine presence or absence of a fever in the ED. If any fever is acceptable, state whether or not time of presentation should be used as the base time (as opposed to time of fever). Finally, abstractors suggested that it be noted whether a statement by the patient or caregiver indicating a fever (temperature greater than 38° C / 100.4° F) obtained prior to arriving at the ED qualifies as fever in the ED. Although our testing results for this measure do not include these changes, they should be considered prior to subsequent implementation of this measure.

Section 13. Summary Statement

Provide a summary rationale for why the measure should be selected for use, taking into account a balance among desirable attributes and limitations of the measure. Highlight specific advantages that this measure has over alternative measures on the same topic that were considered by the measure developer or specific advantages that this measure has over existing measures. If there is any information about this measure that is important for the review process but has not been addressed above, include it here.

This measure, *Appropriate Emergency Department Blood Testing for Children with Sickle Cell Disease*, assesses the percentage of children younger than 18 years of age identified as having SCD presenting to an ED with fever during the measurement year, who 1) received a pulse oximetry reading and 2) had a complete blood count, 3) reticulocyte count, and 4) blood culture performed within 60 minutes following initial contact. This measure uses medical record data to calculate individual rates for the four tests, as well as an overall rate that is a composite of the four individual rates. A higher proportion indicates better performance, as reflected by appropriate testing. There are no existing quality measures for appropriate blood testing in the ED for children with SCD.

Clinical guidelines indicate that all children with SCD who present to the ED with a fever and other signs of infection should be evaluated promptly. Tests should include a pulse oximetry reading, complete blood count, reticulocyte count, and a blood culture. Because the spleen is often compromised at an early age in children with SCD, infection is a frequent and serious complication. Septicemia and meningitis caused by *Streptococcus pneumoniae* are major cause of mortality in children with SCD under the age of 2 years. The presence of fever in children with SCD should be approached with high suspicion for systemic infection. It is crucial that families and clinicians understand that in children with SCD, a temperature of 38 degrees C is an emergency. However, caring for children in the ED is complex from both a medical and behavioral perspective. Children vary developmentally; their ability to communicate and degree of independence may vary widely. For children with SCD, physical challenges and neurocognitive deficits can make ED visits even more complicated. Also, families of children with SCD have reported receiving inappropriate care in the ED stemming from both a lack of familiarity with the children presenting and underdeveloped quality standards for treating children with SCD in the ED.

This measure was tested among a total of 123 children younger than 18 years of age with sickle cell disease. Overall, appropriate blood testing was conducted on 10 percent of children with SCD (range: 7 percent- 10 percent). Pulse oximetry was conducted within 60 minutes of initial contact in the ED for 51 percent of children (range among the three hospitals was: 39 percent-100 percent). Similarly, a complete blood count was conducted within 60 minutes of initial contact in the ED for 25 percent of children (range: 20 percent-26 percent); a reticulocyte count was conducted within 60 minutes of initial contact in the ED for 24 percent of children (range: 20-25 percent); a blood culture was conducted within 60 minutes of initial contact in the ED for 39 percent of children (range: 7 percent-44 percent).

This measure provides families, providers, and purchasers with a straightforward means of assessing how well basic levels of comprehensive care are being provided for children with SCD, including in the ED. The primary information needed for this measure includes basic demographics, dates, diagnostic codes, and procedure codes, all of which are widely available. Continuing advances in the development and implementation of health information technology may establish the feasibility of regularly implementing this measure with data supplied by electronic medical records.

Section 14: Identifying Information for the Measure Submitter

First Name: Gary L.
Last Name: Freed, MD, PhD
Title: Percy and Mary Murphy Professor of Pediatrics, School of Medicine;
Professor of Health Management and Policy, School of Public Health
Organization: University of Michigan
Mailing Address: 300 North Infalls, Room 6E08
City: Ann Arbor
State: MI
Postal Code: 48109
Telephone: 734-615-0616
Email: gfreed@med.umich.edu

The CHIPRA Pediatric Quality Measures Program (PQMP) Candidate Measure Submission Form (CPCF) was approved by the Office of Management and Budget (OMB) in accordance with the Paperwork Reduction Act.

The OMB Control Number is 0935-0205 and the Expiration Date is December 31, 2015.

Public Disclosure Requirements

Each submission must include a written statement agreeing that, should U.S. Department of Health and Human Services accept the measure for the 2014 and/or 2015 Improved Core Measure Sets, full measure specifications for the accepted measure will be subject to public disclosure (e.g., on the Agency for Healthcare Research and Quality [AHRQ] and/or Centers for Medicare & Medicaid Services [CMS] websites), except that potential measure users will not be permitted to use the measure for commercial use. In addition, AHRQ expects that measures and full measure specifications will be made reasonably available to all interested parties. "Full measure specifications" is defined as all information that any potential measure implementer will need to use and analyze the measure, including use and analysis within an electronic health record or other health information technology. As used herein, "commercial use" refers to any sale, license or distribution of a measure for commercial gain, or incorporation of a measure into any product or service that is sold, licensed or distributed for commercial gain, even if there is no actual charge for inclusion of the measure. This statement must be signed by an individual authorized to act for any holder of copyright on each submitted measure or instrument. The authority of the signatory to provide such authorization should be described in the letter.

AHRQ Publication No. 14(17)-P007-8-EF
August 2017